



Pinus strobus 'Glauca' 'Glauca' Eastern White Pine¹

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

This variety of Eastern White Pine has light blue-green needles borne in groups of five (Fig. 1). Unlike the species, needle color is fairly consistent from one tree to the next. The species is the state tree of Maine and Missouri. Although the species can grow more than 100 feet tall with a three to five-foot-diameter trunk and spread 50 to 60 feet, this cultivar is probably going to grow somewhat shorter in landscapes. Growth is very rapid at first but slows down with age.

GENERAL INFORMATION

Scientific name: *Pinus strobus* 'Glauca'

Pronunciation: PIE-nus STROE-bus

Common name(s): 'Glauca' Eastern White Pine

Family: *Pinaceae*

USDA hardiness zones: 3B through 7 (Fig. 2)

Origin: native to North America

Uses: Bonsai; hedge; recommended for buffer strips around parking lots or for median strip plantings in the highway; screen; shade tree; specimen; Christmas tree; no proven urban tolerance

Availability: grown in small quantities by a small number of nurseries

DESCRIPTION

Height: 50 to 80 feet

Spread: 50 to 60 feet

Crown uniformity: symmetrical canopy with a regular (or smooth) outline, and individuals have more or less identical crown forms

Crown shape: oval; pyramidal




Figure 1. Middle-aged 'Glauca' Eastern White Pine.

Crown density: moderate

Growth rate: medium

Texture: fine

Foliage

Leaf arrangement: alternate; spiral (Fig. 3)

Leaf type: simple

Leaf margin: entire

Leaf shape: needle-like (filiform)

1. This document is adapted from Fact Sheet ST-476, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: October 1994.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

Leaf venation: parallel

Leaf type and persistence: evergreen; fragrant;
needle leaf evergreen

Leaf blade length: 2 to 4 inches

Leaf color: blue or blue-green

Fall color: no fall color change

Fall characteristic: not showy

Flower

Flower color: pink; yellow

Flower characteristics: inconspicuous and not
showy; spring flowering

Fruit

Fruit shape: elongated

Fruit length: 6 to 12 inches

Fruit covering: dry or hard

Fruit color: brown

Fruit characteristics: does not attract wildlife; fruit,
twigs, or foliage cause significant litter; persistent on
the tree; showy

Trunk and Branches

Trunk/bark/branches: bark is thin and easily
damaged from mechanical impact; droop as the tree
grows, and will require pruning for vehicular or
pedestrian clearance beneath the canopy; showy trunk;
should be grown with a single leader; no thorns

Pruning requirement: needs little pruning to develop
a strong structure

Breakage: susceptible to breakage either at the crotch
due to poor collar formation, or the wood itself is
weak and tends to break

Current year twig color: brown; green

Current year twig thickness: thin

Wood specific gravity: 0.35

Culture

Light requirement: tree grows in part shade/part sun;
tree grows in full sun

Soil tolerances: loam; sand; acidic; well-drained

Drought tolerance: moderate

Aerosol salt tolerance: none




Figure 3. Foliage of 'Glauca' Eastern White Pine.

Other

Roots: surface roots are usually not a problem

Winter interest: no special winter interest

Outstanding tree: not particularly outstanding

Invasive potential: little, if any, potential at this time

Verticillium wilt susceptibility: not known to be susceptible

Pest resistance: very sensitive to one or more pests or diseases which can affect tree health or aesthetics

USE AND MANAGEMENT

Several branches on young trees normally originate from the same point on the trunk forming a tree appearing to be built of layers of foliage. Although young trees are pyramidal and usually grow with one central leader, the layers (or whorls) of horizontal branches give White Pine a distinctive appearance in middle and old age. The gray bark on the trunk and large branches remains unusually smooth through middle age, breaking up into elongated blocks in old age. Be sure to purchase only certified rust-resistant plants.

The lower branches are retained making White Pine an excellent candidate for specimen use, although group plantings with trees spaced 15 to 25 feet apart add a soft accent to any landscape. Planted 8 to 15 feet apart they are one of only a few Pines which makes a nice hedge or screen of soft foliage. They are also used as a Christmas tree. Pinch or clip the candles as the needles begin to emerge.

Root systems are usually shallow and highly branched with many fine roots close to the surface of the soil. Trees transplant well balled and burlapped or from containers. The wood is soft and is popular as a lumber tree in the northern part of the country.

Young White Pines are quite tolerant of half-day shade while mature White Pines prefer a sunny location and tolerate loamy, moist, well-drained soils. They do not grow well and often die on clay or on soil with a pH above 7. Trees appear to have little tolerance for drought, soil compaction and heat and should be used only in the cooler climates. Eastern White Pine is susceptible to salt injury from roads or drain fields and is sensitive to air pollution (particularly ozone and sulfur dioxide).

There are a few other cultivars: 'Fastigiata' - nearly columnar habit; 'Nana' - a dwarf, compact with short needles; 'Pendula' - weeping; 'Prostrata' - absolutely prostrate.

Pests

White Pine weevil is probably the biggest problem. The larvae of White Pine weevils feed on the sapwood of the leaders. The leader is killed and the multiple shoots replacing it form a bushy head of needles. First symptoms are pearl white drops of resin on the leaders. The leaders die when the shoot is girdled as adults emerge in summer.

Some adelgids will appear as white cottony growths on the bark. All types produce honeydew which may support sooty mold. European Pine shoot moth causes young shoots to fall over. Infested shoots may exude resin. The insects can be found in the shoots during May.

Bark beetles bore into trunks making small holes scattered up and down the trunk. The holes look like shotholes. Stressed trees are more susceptible to attack. Keep trees healthy.

Sawfly larvae caterpillars are variously colored but generally feed in groups on the needles. Some sawfly larvae will flex or rear back in unison when disturbed. Sawflies can cause rapid defoliation of branches if left unchecked.

Pine needle miner larvae feed inside needles causing them to turn yellow and dry up.

Pine needle scale is a white, elongated scale found on the needles. Pine tortoise scale is brown and found on twigs. Depending on the scale, horticultural oil may control overwintering stages.

Pine spittle bug lives and hides in a foamy mass.

Spruce mites cause damage to older needles, and are usually active in the spring and fall. Mites cause older needles to become yellowed or stippled.

Zimmerman Pine moth larvae bore into the trunk. The only outward symptoms may be death of parts of the tree or masses of hardened pitch on the branches.

Diseases

Procerva root rot kills most White Pines planted off site. Avoid planting in dry sites and never plant in clay or alkaline soil.

White Pine blister rust attacks White Pine and uses currant as an alternate host. European Black Currant, the favored alternate host, may be banned from certain areas. Other Currants, particularly Red Currant should not be grown within 300 feet of Pines. Infected branches may be pruned off. Be sure to select White Pine trees certified to be rust-resistant.

Canker diseases may rarely cause dieback of landscape Pines. Keep trees healthy and prune out the infected branches.

Needle cast is common on small trees and plantation or forest trees. Infected needles yellow and fall off.

White Pine decline is used to describe the slow decline of trees planted in dry, clay soils low in organic matter. Plants with this disorder have only a small cluster of needles at the ends of the branches.