Aesculus parviflora1

Edward F. Gilman²

Introduction

Native from Alabama and South Carolina into Florida, Bottlebrush Buckeye forms a rounded mass of dark green, palmately-compound foliage in mid-spring (Fig. 1). The shrub eventually reaches about 8-feet-tall but grows to 12-feet-wide. It can be found in its native, moist, shaded habitat flowering in early summer. The delicate, showy, white flowers are held well above the foliage in terminal panicles up to 12 inches long. Bottlebrush Buckeye has been successfully used as far north as Chicago (hardiness zone 5).

General Information

Scientific name: Aesculus parviflora

Pronunciation: ESS-kew-lus par-vif-FLOR-uh **Common name(s):** Bottlebrush Buckeye

Family: Hippocastanaceae

Plant type: shrub

USDA hardiness zones: 5 through 9A (Fig. 2) **Planting month for zone 7:** year round **Planting month for zone 8:** year round **Planting month for zone 9:** year round

Origin: native to Florida

Uses: specimen; screen; foundation; border

Availablity: somewhat available, may have to go out of the

region to find the plant

Description

Height: 5 to 10 feet **Spread:** 10 to 15 feet



Figure 1. Bottlebrush Buckeye.

Plant habit: round Plant density: moderate Growth rate: slow Texture: coarse

Foliage

Leaf arrangement: opposite/subopposite

- This document is Fact Sheet FPS-17, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: October 1999. Please visit the EDIS web site at http://edis.ifas.ufl.edu.
- Edward F. Gilman, professor, Environmental Horticulture Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University
 of Florida, Gainesville, 32611.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service / Institute of Food and Agricultural Sciences / University of Florida / Christine Taylor Waddill, Dean




Figure 2. Shaded area represents potential planting range.

Leaf type: palmately compound

Leaf margin: crenate
Leaf shape: oblong; obovate
Leaf venation: pinnate

Leaf type and persistence: deciduous **Leaf blade length:** 4 to 8 inches

Leaf color: green **Fall color:** yellow

Fall characteristic: showy

Flower

Flower color: white

Flower characteristic: spring flowering

Fruit

Fruit shape: elongated Fruit length: 1 to 3 inches Fruit cover: dry or hard Fruit color: brown

Fruit characteristic: showy

Trunk and Branches

Trunk/bark/branches: typically multi-trunked or clumping

stems; not particularly showy

Current year stem/twig color: gray/silver Current year stem/twig thickness: thick

Culture

Light requirement: plant grows in part shade/part sun; plant

grows in the shade

Soil tolerances: extended flooding; acidic; sand; loam; clay;

Drought tolerance: moderate **Soil salt tolerances:** poor **Plant spacing:** 36 to 60 inches

Other

Roots: sprouts from roots or lower trunk **Winter interest:** no special winter interest

Outstanding plant: plant has outstanding ornamental features

and could be planted more

Invasive potential: not known to be invasive

Pest resistance: long-term health usually not affected by pests

Use and Management

Allow plenty of room for this spreading shrub since it looks best without pruning. Pruning ruins the natural uniform shape. Locate it in the partial or full shade for a splash of color in early summer. Fall color is yellow, occasionally developing into a short-lived showy display.

Pests and Diseases

Few problems are reported on this nice, native plant.




Figure 3. Foliage of Bottlebrush Buckeye