



Cooperative Extension Service
Institute of Food and Agricultural Sciences

***Antirrhinum majus* 'Liberty'**¹

Edward F. Gilman²

Introduction

These Snapdragon selections grow no more than about 2-foot-tall (Fig. 1). The flowers come in a wide range of colors from reds, orange, yellow, and maroon. Plants with dark colored flowers have dark green or reddish stems and those with white or pale flowers have pale green stems.

General Information

Scientific name: *Antirrhinum majus* 'Liberty'

Pronunciation: an-tur-RYE-num MAY-jus

Common name(s): 'Liberty' Snapdragon

Family: *Scrophulariaceae*

Plant type: annual

USDA hardiness zones: all zones (Fig. 2)

Planting month for zone 7: Apr; Sep

Planting month for zone 8: Feb; Mar; Oct; Nov; Dec

Planting month for zone 9: Feb; Oct; Nov; Dec

Planting month for zone 10 and 11: Feb; Nov; Dec

Origin: not native to North America

Uses: mass planting; container or above-ground planter; cut flowers; edging

Availability: generally available in many areas within its hardiness range

Description

Height: 1 to 2 feet

Spread: .5 to 1 feet

Plant habit: upright

Plant density: moderate




Figure 1. 'Liberty' Snapdragon.

Growth rate: fast

Texture: medium

Foliage

Leaf arrangement: alternate

Leaf type: simple

Leaf margin: entire

1. This document is Fact Sheet FPS-45, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: October 1999. Please visit the EDIS web site at <http://edis.ifas.ufl.edu>.
2. Edward F. Gilman, professor, Environmental Horticulture Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service / Institute of Food and Agricultural Sciences / University of Florida / Christine Taylor Waddill, Dean



Figure 2. Shaded area represents potential planting range.

Leaf shape: oblong; spatulate
Leaf venation: none, or difficult to see
Leaf type and persistence: not applicable
Leaf blade length: 2 to 4 inches
Leaf color: green
Fall color: not applicable
Fall characteristic: not applicable

Flower

Flower color: white; orange; lavender
Flower characteristic: showy

Fruit

Fruit shape: no fruit
Fruit length: no fruit
Fruit cover: no fruit
Fruit color: not applicable
Fruit characteristic: inconspicuous and not showy

Trunk and Branches

Trunk/bark/branches: not applicable
Current year stem/twig color: green

Current year stem/twig thickness: thick

Culture

Light requirement: plant grows in part shade/part sun
Soil tolerances: acidic; slightly alkaline; clay; sand; loam
Drought tolerance:
Soil salt tolerances: unknown
Plant spacing: 6 to 12 inches

Other

Roots: not applicable
Winter interest: not applicable
Outstanding plant: not particularly outstanding
Invasive potential: not known to be invasive
Pest resistance: long-term health usually not affected by pests

Use and Management

Snapdragons grow in any slightly acid, garden soil, however, they don't grow well in unamended clay. The plants require full sun and moist soil. A second crop of flowers may be obtained from plants that have finished flowering. Cut them back to within 5 or 6 nodes of the ground when the first flowers fade. Fertilize when the second crop of flower buds become visible.

Snapdragons may be propagated by seeds, or by cuttings which root readily. The seed germinates in 10 to 14 days at 70-degrees F. Do not cover the seed with soil. Prechilled seeds germinate best. Seedlings with two to three sets of leaves are pinched. Set plants in the ground after the danger of frost has passed. Plant in the fall for winter color in USDA hardiness zones 9 to 11. Plants sometimes survive and flower throughout the winter in zone 8b. Set the plants six to ten inches apart.

Aphids feed on terminal growth and the underside of the leaves. The insects suck juices, and heavy infestations seriously weaken the plants.

The greenhouse leaf tier chews irregular-shaped areas in the leaves and webs the leaves together. Pesticides are seldom effective after the insect rolls the leaves.

Mites cause a bronzed or stippled appearance on the foliage, especially in hot weather.

Pests and Diseases

Rust causes brown pustules surrounded by yellowed tissue on the leaves. Plants may bloom prematurely, have small flowers, and die early. Use proper plant spacings and resistant varieties.

Anthracnose attacks the leaves and stems in late summer. On older stems the spots are sunken, oblong, yellowish-green to gray with a narrow brown border. On the leaves, the spots are yellowish green turning dirty white with a narrow brown border. When the stem is girdled the plant dies. Destroy infected plants and use wider spacings.

Gray mold causes flower spikes to wilt and light brown areas form on the lower stem of the flower cluster. Infected plants break over below the flowers. The disease is worse in wet weather. Cut off infected flower stalks and keep beds free of debris.

Stem rot can be detected by the presence of cottony growth on stems of infected plants near the soil line. Infected plants die and should be destroyed.