



Cooperative Extension Service  
Institute of Food and Agricultural Sciences

## *Narcissus spp.*<sup>1</sup>

Edward F. Gilman<sup>2</sup>

### Introduction

The well-known Daffodil is now available in single or double flower forms in various color combinations of yellow, white, apricot, cream, peach, red, or orange (Fig. 1). Long used in forced pot culture, Daffodils make attractive naturalized groundcovers in sweeping drifts or under trees. Daffodils also work well under shrubs, in rock gardens or near water, or used as an edging plant. Few plants seem to signify the freshness of spring quite as well as Daffodils.

### General Information

**Scientific name:** *Narcissus spp.*

**Pronunciation:** nar-SIS-suss species

**Common name(s):** Daffodil, Narcissus

**Family:** *Amaryllidaceae*

**Plant type:** perennial; herbaceous

**USDA hardiness zones:** 3B through 10 (Fig. 2)

**Planting month for zone 7:** year round

**Planting month for zone 8:** year round

**Planting month for zone 9:** year round

**Planting month for zone 10:** year round

**Origin:** not native to North America

**Uses:** mass planting; border; container or above-ground planter; edging; suitable for growing indoors

**Availability:** generally available in many areas within its hardiness range



Figure 1. Daffodil.

**Spread:** .5 to 2 feet

**Plant habit:** upright

**Plant density:** moderate

**Growth rate:** moderate

**Texture:** fine

### Description

**Height:** 1 to 2 feet

1. This document is Fact Sheet FPS-423, one of a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: October, 1999 Please visit the EDIS Web site at <http://edis.ifas.ufl.edu>.
2. Edward F. Gilman, professor, Environmental Horticulture Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

The Institute of Food and Agricultural Sciences is an equal opportunity/affirmative action employer authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, sex, age, handicap, or national origin. For information on obtaining other extension publications, contact your county Cooperative Extension Service office. Florida Cooperative Extension Service / Institute of Food and Agricultural Sciences / University of Florida / Christine Taylor Waddill, Dean



Figure 2. Shaded area represents potential planting range.

### Foliage

**Leaf arrangement:** most emerge from the soil, usually without a stem

**Leaf type:** simple

**Leaf margin:** entire

**Leaf shape:** linear

**Leaf venation:** parallel

**Leaf type and persistence:** deciduous

**Leaf blade length:** 12 to 18 inches

**Leaf color:** green

**Fall color:** no fall color change

**Fall characteristic:** not showy

### Flower

**Flower color:** white; yellow, cream; apricot; red; orange

**Flower characteristic:** spring flowering; pleasant fragrance; winter flowering

### Fruit

**Fruit shape:** unknown

**Fruit length:** unknown

**Fruit cover:** unknown

**Fruit color:** green

**Fruit characteristic:** inconspicuous and not showy

### Trunk and Branches

**Trunk/bark/branches:** not applicable

**Current year stem/twig color:** not applicable

**Current year stem/twig thickness:** not applicable

## Culture

**Light requirement:** plant grows in part shade/part sun

**Soil tolerances:** occasionally wet; acidic; sand; loam; clay

**Drought tolerance:** high

**Soil salt tolerances:** poor

**Plant spacing:** 6 to 12 inches

## Other

**Roots:** not applicable

**Winter interest:** plant has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

**Outstanding plant:** not particularly outstanding

**Invasive potential:** not known to be invasive

**Pest resistance:** long-term health usually not affected by pests

## Use and Management

Hundreds of cultivars exist today and planting dates will vary accordingly, but the bulbs are usually planted in fall when the soil is cool. Although Daffodils grow well in full sun or light shade, the flowers last longer in light shade. The bulbs are planted with four to five inches of soil on top of them. If spaced eight inches apart, the bulbs should not need to be divided for two to three years. Double-nosed bulbs are actually two blooming sized bulbs joined together. When selecting a location for planting, it should be noted that the individual flowers will face the sun.

Bulbs should be watered well after planting and, in the south, thereafter if winters are dry. After flowering, plants should continue to be watered and fertilized. The foliage should be allowed to naturally turn yellow and die without being removed as this helps supply energy to the bulb for the next year's bloom. When flowering becomes sparse and the blooms small, division of the bulbs may be necessary. After the foliage has died down, dig and lift the bulbs, separating off only those bulbs which come off easily. Replant at once or store at 45-degrees F. for a very short period.

The tulip bulb aphid damages stored bulbs and feeds on the leaves and other above ground parts in the spring.

The bulb mite causes infested bulbs to produce stunted leaves and deformed flowers.

Bulb flies will infest Narcissus. The larvae make the bulbs soft and unable to grow. The openings created by the insect are entrances for diseases.

Millipedes will feed on the undersides of bulbs.

The stem and bulb nematode causes darkened bulb scales giving a ringed appearance to bulb cross sections. Yellowish pockets in the bulb contain many nematodes. Infested bulbs do not grow, or fail to flower. The shoots are abnormal and twisted. Remove infested plants and avoid rich wet soil with a high humus content.

## Pests and Diseases

Blue mold is a storage problem and is worse on injured bulbs. The disease is most likely to occur in moist storage conditions.

Crown rot covers bulbs with layers of coarse white mold. Remove infected plants and avoid infested soil for two to three years.

Basal rot causes a decay starting at the roots or bases of the scales, and spreads upwards inside the bulb. The plants are dwarfed and the blossoms abnormal.

Fire causes spotting and rotting of flowers in humid weather. The disease spreads to, and rapidly destroys, the leaves. The symptoms on leaves are dark reddish, elongated, brown spots. Rake up and destroy old foliage.

Leaf spot causes large spots or blotches on the leaves. The leaves wither and die. The bulbs lack nourishment due to the premature death of leaves. Remove and destroy plants with diseased foliage.

Smolder rots the foliage and flowers in cold, wet seasons. The leaves are stuck together when they emerge and infected bulbs rot in storage. Remove and destroy infected plants.

Root rot causes leaf yellowing and stunting and withering of plants. The roots have yellowish or brown discolored stripes and decayed regions. The bulbs remain sound. This disease is more of a problem where narcissi have been planted for several years.

Leaf scorch causes blighted leaf tips to be separated from healthy portions by a yellow area. The secondary infection on the lower leaves is minute, water-soaked or yellowish spots, which become raised, scabby and reddish. The flower stalks and flowers may be spotted.