



Gleditsia triacanthos var. *inermis* Thornless Honeylocust¹

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

Honeylocust grows quickly to 70 feet or more with an oval or rounded canopy (Fig. 1). The species has undesirable thorns on the trunk and main branches and large seed pods so it is best to plant selections of the variety *inermis* which are both thornless and some nearly seedless. Some leaves on the tree are bipinnately compound, others are pinnately compound. The seed pods look rather unsightly hanging on the tree into the fall and make quite a mess as they litter the ground below the canopy. The tree is strong-wooded and casts light shade. Lawns grow fairly well beneath the tree and there is little to rake up in the fall since the tiny leaflets filter in between the blades of grass or are washed away in the rain. Honeylocust has a yellow or golden fall color in the northern part of its range. Trees often defoliate early in the south and are bare by October.

GENERAL INFORMATION

Scientific name: *Gleditsia triacanthos* var. *inermis*

Pronunciation: gleh-DIT-see-uh try-uh-KANTH-oase
variety ih-NER-miss

Common name(s): Thornless Honeylocust

Family: *Leguminosae*

USDA hardiness zones: 3 through 8A (Fig. 2)

Origin: native to North America

Uses: large parking lot islands (> 200 square feet in size); wide tree lawns (>6 feet wide); recommended for buffer strips around parking lots or for median strip plantings in the highway; reclamation plant; shade tree; specimen; sidewalk cutout (tree pit); residential street tree; tree has been successfully grown in urban




Figure 1. Young Thornless Honeylocust.

areas where air pollution, poor drainage, compacted soil, and/or drought are common

Availability: generally available in many areas within its hardiness range

DESCRIPTION

Height: 50 to 75 feet

Spread: 35 to 50 feet

Crown uniformity: irregular outline or silhouette

1. This document is adapted from Fact Sheet ST-279, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

Crown shape: oval; round

Crown density: open

Growth rate: fast

Texture: fine

Foliage

Leaf arrangement: alternate (Fig. 3)

Leaf type: bipinnately compound; odd pinnately compound

Leaflet margin: crenate

Leaflet shape: lanceolate; oblong

Leaflet venation: pinnate

Leaf type and persistence: deciduous

Leaflet blade length: less than 2 inches

Leaf color: green

Fall color: copper; yellow

Fall characteristic: showy

Flower

Flower color: yellow

Flower characteristics: pleasant fragrance; inconspicuous and not showy; spring flowering

Fruit

Fruit shape: elongated; pod

Fruit length: 12 inches or more; 6 to 12 inches

Fruit covering: dry or hard

Fruit color: brown; purple

Fruit characteristics: does not attract wildlife; fruit, twigs, or foliage cause significant litter; persistent on the tree; showy

Trunk and Branches

Trunk/bark/branches: grow mostly upright and will not droop; showy trunk; should be grown with a single leader; no thorns

Pruning requirement: requires pruning to develop strong structure

Breakage: resistant

Current year twig color: brown

Current year twig thickness: thick

Culture

Light requirement: tree grows in part shade/part sun; tree grows in full sun




Figure 3. Foliage of Thornless Honeylocust.

Soil tolerances: clay; loam; sand; acidic; occasionally wet; alkaline; well-drained

Drought tolerance: high

Aerosol salt tolerance: high

Soil salt tolerance: good

Other

Roots: surface roots can lift sidewalks or interfere with mowing

Winter interest: tree has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

Outstanding tree: not particularly outstanding

Invasive potential: little, if any, potential at this time

Ozone sensitivity: sensitive or moderately tolerant

Verticillium wilt susceptibility: not known to be susceptible

Pest resistance: long-term health usually not affected by pests

USE AND MANAGEMENT

Some cultivars, especially 'Skyline' grow a central leader and would require little pruning but others grow many upright codominant trunks. These trees will need to be trained when they are young by two or three prunings spaced several years apart. Strive to develop one central trunk with upright spreading branches spaced several feet apart along the trunk. Purchase good quality trees with one leader in order to reduce the pruning requirement.

The tree has no particular soil preferences and is useful in dry or alkaline areas, although its native habitat is along stream banks. It tolerates compacted, poorly aerated soil and flooding for a period of time and does well in confined soil spaces. Honeylocust adapts well as a city street tree and is tolerant to small planting pits in concrete. It is susceptible to breakage in ice storms.

Unfortunately, it has been overplanted in some areas and insect problems are beginning to catch up with Honeylocust, including the cultivars. Recommend planting in moderation to avoid catastrophe if insects or diseases invade. It might be best to plant Pistacia, Zelkova, Taxodium, Quercus or some other proven urban tough tree in place of Honeylocust to avoid potential insect, disease and early defoliation problems in the South.

Most garden centers will have at least one cultivar of Honeylocust in stock. Some of the cultivars may develop thorns and/or seed pods when they get older and they may be best suited for areas north of USDA hardiness zone 8b. The cultivars are: 'Cottage Green' - semi upright, seedless, thornless; 'Imperial' - upright-spreading, seedless, and thornless until 10 to 15-years-

old when some seeds do develop; 'Majestic' - upright, seedless, thornless; 'Maxwell' - upright, seedless, thornless; 'Moraine' - spreading, usually seedless, thornless; 'Rubylace' - new reddish foliage, seedless, thornless, color not outstanding, may need staking when young; 'Shademaster' - upright, spreading, usually seedless and thornless until 10 to 15-years-old when some seeds do develop - perhaps the best cultivar; 'Skyline' - pyramidal with a dominant central leader, generally seedless, thornless; 'Sunburst' - new yellow foliage, seedless, thornless, favored by plant bugs and leafhoppers.

There is a leaf spot may be a problem. Rake up and dispose of infected leaves.

Powdery mildew may cause a white coating on the leaves but is seldom serious.

Pests

Mimosa webworm has become a serious pest on Honeylocust in some communities.

Boring insects may be largely prevented by keeping trees healthy with regular fertilization. They usually attack trees under stress from other problems.

The combination of plant bug and leafhopper feeding causes the leaves to drop. Plant bugs may be more common on the yellow leaved cultivar 'Sunburst' than on green leaved types. Both insects are green so they will be hard to detect.

Pod gall midge causes unusual reddish galls at the tips of the branches. Leaflets become pod-like. The galls appear in late spring and may be most common on thornless, seedless cultivars. These have become quite a problem in many areas. Control is difficult.

Spider mites cause an autumn-like yellowing of the leaves. Diagnosis of this problem is difficult due to the small size of the insect and leaflets. Look for the mites and their webbing near the midrib at the base of the leaflets.

Leafminers and bagworm can also be a problem.

Aphid infestations can be troublesome.

Diseases

Canker causing fungi or bacteria attack branches and trunks causing dieback of parts or the entire tree. Keep the trees healthy and avoid unnecessary wounding. Infected areas have discolored bark, peeling bark, discolored sapwood, or a crack between the diseased and healthy bark. The *ronectria* canker is especially damaging.