

## *Liquidambar styraciflua* 'Burgundy' 'Burgundy' Sweetgum<sup>1</sup>

Edward F. Gilman and Dennis G. Watson<sup>2</sup>

### INTRODUCTION

This cultivar of Sweetgum grows in a narrow pyramid to a height of 75 feet and may spread to 50 feet (Fig. 1). The beautifully glossy, star-shaped leaves turn bright red-purple in the fall (USDA hardiness zones 6 and 7) and early winter (USDA hardiness zones 8 and 9). In the fall, leaves are held on the tree longer than the species. It is well adapted to the Deep South. On some trees, particularly in the northern part of its range, branches are covered with characteristic corky projections. The trunk is normally straight and does not divide into double or multiple leaders and side branches are small in diameter on young trees, creating a pyramidal form. The bark becomes deeply ridged at about 25-years-old. Sweetgum makes a nice conical park, campus or residential shade tree for large properties when it is young, developing a more oval or rounded canopy as it grows older as several branches become dominant and grow in diameter.

### GENERAL INFORMATION

**Scientific name:** *Liquidambar styraciflua* 'Burgundy'

**Pronunciation:** lick-wid-AM-bar  
sty-rass-ih-FLOO-uh

**Common name(s):** 'Burgundy' Sweetgum

**Family:** *Hamamelidaceae*

**USDA hardiness zones:** 6 through 10A (Fig. 2)

**Origin:** native to North America

**Uses:** large parking lot islands (> 200 square feet in size); wide tree lawns (>6 feet wide); recommended for buffer strips around parking lots or for median strip plantings in the highway; reclamation plant; shade




Figure 1. Young 'Burgundy' Sweetgum.

tree; specimen; residential street tree; no proven urban tolerance

**Availability:** somewhat available, may have to go out of the region to find the tree

1. This document is adapted from Fact Sheet ST-359, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

## DESCRIPTION

**Height:** 60 to 75 feet  
**Spread:** 35 to 50 feet  
**Crown uniformity:** symmetrical canopy with a regular (or smooth) outline, and individuals have more or less identical crown forms  
**Crown shape:** oval; pyramidal  
**Crown density:** dense  
**Growth rate:** medium  
**Texture:** coarse

## Foliage

**Leaf arrangement:** alternate (Fig. 3)  
**Leaf type:** simple  
**Leaf margin:** lobed; serrate  
**Leaf shape:** star-shaped  
**Leaf venation:** palmate  
**Leaf type and persistence:** deciduous  
**Leaf blade length:** 4 to 8 inches  
**Leaf color:** green  
**Fall color:** purple; red  
**Fall characteristic:** showy

## Flower

**Flower color:** green; yellow  
**Flower characteristics:** inconspicuous and not showy; spring flowering

## Fruit

**Fruit shape:** round  
**Fruit length:** 1 to 3 inches  
**Fruit covering:** dry or hard  
**Fruit color:** brown  
**Fruit characteristics:** attracts birds; attracts squirrels and other mammals; fruit, twigs, or foliage cause significant litter; persistent on the tree; showy

## Trunk and Branches

**Trunk/bark/branches:** droop as the tree grows, and will require pruning for vehicular or pedestrian clearance beneath the canopy; not particularly showy; should be grown with a single leader; no thorns  
**Pruning requirement:** needs little pruning to develop a strong structure  
**Breakage:** resistant  
**Current year twig color:** brown; reddish  
**Current year twig thickness:** medium




Figure 3. Foliage of 'Burgundy' Sweetgum.

**Wood specific gravity:** 0.52

### Culture

**Light requirement:** tree grows in part shade/part sun; tree grows in full sun

**Soil tolerances:** clay; loam; sand; slightly alkaline; acidic; extended flooding; well-drained

**Drought tolerance:** moderate

**Aerosol salt tolerance:** moderate

### Other

**Roots:** surface roots can lift sidewalks or interfere with mowing

**Winter interest:** tree has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

**Outstanding tree:** tree has outstanding ornamental features and could be planted more

**Invasive potential:** seeds itself into the landscape

**Verticillium wilt susceptibility:** not known to be susceptible

**Pest resistance:** long-term health usually not affected by pests

## USE AND MANAGEMENT

Be careful when locating Sweetgum as a street tree since its large, aggressive roots may lift curbs and sidewalks. Plant trees 8 to 10 feet or more from curbs. Some communities have large numbers of Sweetgum planted as street trees. Much of the root system is shallow (particularly in its native, moist habitat), but there are deep vertical roots directly beneath the trunk in well-drained and in some other soils. The fruit may be a litter nuisance to some in the fall, but this is usually only noticeable on hard surfaces, such as roads, patios, and sidewalks, where people could slip and fall on the fruit. The cultivar 'Rotundiloba' is fruitless. The tree should be planted only in soil with a pH of 7 or less. The seeds provide food for wildlife and will often readily germinate in shrub and groundcover beds, requiring their removal to maintain a neat landscape appearance.

Although it grows at a moderate pace, Sweetgum is rarely attacked by pests, and tolerates wet soils, but chlorosis is often seen in alkaline soils. Trees grow well in deep soil, poorly in shallow, droughty soil. Sweetgum is difficult to transplant and should be planted from containers or transplanted in the spring when young since it develops deep roots on well-drained soil. It is native to bottomlands and moist soils and tolerates only some (if any) drought. Existing trees often die-back near the top of the crown, apparently due to extreme sensitivity to construction injury to the root system, or drought injury. The tree leafs out early in the spring and is sometimes damaged by frost.

Cultivars have been selected for their fall color, leaf shape, or growth habit: 'Festival' - narrow upright growth habit, peach-colored fall foliage, less cold hardy, more adapted to the southern part of the range; 'Moraine' is reputed to be the most cold hardy; 'Palo Alto' - pyramidal, symmetrical growth, bright orange fall color; 'Rotundiloba' - round leaf tips, no fruit production, narrow pyramidal form. *Liquidambar formosana* has a broader spreading crown.

### Pests

Bagworm makes sacks by webbing together pieces of leaves. The insects live in the sacks while they feed. Small numbers of insects may be picked off by hand. I

Fall webworm webs over portions of large branches or may completely cover small branches.

The insects feed on leaves inside the nest. If practical, nests can be pruned out while small and when the insects are inside. A few nests in large trees are not serious.

Leaf miner causes brown blotches on leaves. If injury is caused by leaf miner the browned upper and lower leaf surfaces will be completely separate when the leaf is torn in two.

Cottony-cushion scale, Sweetgum scale, and walnut scale can infest the branches. Use horticultural oil to provide some control.

Tent caterpillars make nests to live in but leave the nests to feed. Prune out nests at the tips of small branches. Do not burn the nest while it is still in the tree because you will injure the tree.

## **Diseases**

Sweetgum may be attacked by canker diseases. These diseases cause sunken areas on the trunk and some cause profuse "bleeding". Infected bark and sapwood will be brown and dead. There is no chemical control for canker diseases. Severely infected trees will die. Prune cankers out of lightly infected trees. Maintain tree health by watering and fertilizing.

Leaf spots of various types may attack Sweetgum, causing premature defoliation, but are not serious. Rake up and destroy infected leaves to help control if there are no adjacent Sweetgum to add inoculum.